











- Cozannet P., Yacoubi N., Dumont T., Preynat A., 2019. Une nouvelle génération de multi-carbohydase riche en xylanase et arabinofuranosidase efficace pour améliorer la digestibilité des aliments de porcs. Journées Rech. Porcine 51, 93-98.
- Cromwell G.L., Stahly T.S., Randolph J.H., 1991. Effects of phytase on the utilization of phosphorus in corn-soybean meal diets by growing-finishing pigs. *J. Anim. Sci.*, 69, 358 (Abstr. 1.).
- Ekpe E.D., Zijlstra R.T., Patience J.F., 2002. Digestible phosphorus requirement of grower pigs. *Can. J. Anim. Sci.*, 82, 541-549.
- Gloaguen M., Le Floch N., Brossard L., Barea R., Primot Y., Corrent E., van Milgen J., 2012. Response of piglets to the valine content in the diet in combination with the supply of other branched-chain amino acids. *Animal*, 5, 1734-1742.
- Harper A.F., Kornegay E.T., Schell T.C., 1997. Phytase supplementation of low-phosphorus growing-finishing pig diets improves performance, phosphorus digestibility, and bone mineralization and reduces phosphorus excretion. *J. Anim. Sci.*, 75, 3174-3186.
- Lawlor P.G., Cozannet P., Ryan W.F., Lynch P.B., 2019. Effect of a combination phytase and multi-carbohydrolase enzyme supplement on growth performance and bone mineralization of pigs from six weeks to slaughter at 105 kg. *Livest. Sci.*, 223, 144-150.
- Lindberg J.E., Lyberg K., Sands J., 2007. Influence of phytase and xylanase supplementation of a wheat-based diet on ileal and total tract digestibility in growing pigs. *Livest. Sci.*, 109, 268-270.
- Liu K.G., Bello A., Jlali M., Cozannet P., Wu D., Davin R., Preynat A., 2019. Evaluation of the efficacy of a Multi carbohydase and phytase on corn-wheat-soybean meal-based diets with varying levels in metabolizable energy, digestible amino acids, available phosphorus and calcium. *Proceeding Australian Poultry Science Symposium.*, 30<sup>th</sup>, 172-176.
- Liu S.Y., Naranjo V.D., Chrystal P.V., Buyse J., Selle P.H., 2017. Box-Behnken optimisation of growth performance, plasma metabolites and carcass traits as influenced by dietary energy, amino acid and starch to lipid ratios in broiler chickens. *Plos One*, 14, e0213875.
- Moehn S., Atakora J.K.A., Sands J., Ball R.O., 2007. Effect of phytase-xylanase supplementation to wheat-based diets on energy metabolism in growing-finishing pigs fed ad libitum. *Livest. Sci.*, 109, 271-274.
- Nortey T.N., Patience J.F., Simmins P.H., Trottier N.L., Zijlstra R.T., 2007. Effects of individual or combined xylanase and phytase supplementation on energy, amino acid, and phosphorus digestibility and growth performance of grower pigs fed wheat-based diets containing wheat millrun. *J. Anim. Sci.*, 85, 1432-1443.
- NRC. 2012. Nutrient requirements of swine. National Academic Press, Washington, DC. nb pp?
- Nyachoti C.M., Zijlstra R.T., de Lange C.F.M., Patience J.F., 2004. Voluntary feed intake in growing-finishing pigs: A review of the main determining factors and potential approaches for accurate predictions. *Can. J. Anim. Sci.*, 84, 549-566.
- Rosenfelder-Kuon P., Siegert W., Rodehutschord M., 2019. Effect of microbial phytase supplementation on P digestibility in pigs: a meta-analysis. *Arch. Anim. Nut.*, 74, 1-18.
- Sauber T.E., Owens F.N., 2001. Cereal grains and byproducts for swine. *Swine Nutrition*. A. J. Lewis and L. L. Southern (ed), New York, CRC Press, 785-802.
- Sharma N.K., Choct M., Toghyani M., Laurenson Y.C.S.M., Giriash C.K., Swick R.A., 2018. Dietary energy, digestible lysine, and available phosphorus levels affect growth performance, carcass traits, and amino acid digestibility of broilers. *Poult. Sci.* 97, 1189-1198.
- Shaw D.T., Rozeboom D.W., Hil G.M., Orth M.W., Rosenstein D.S., Link J.E., 2006. Impact of supplement withdrawal and wheat middling inclusion on bone metabolism, bone strength, and the incidence of bone fractures occurring at slaughter in pigs. *J. Anim. Sci.*, 84, 1138-1146.
- Quiniou N., Noblet J., 2012. Effect of the dietary net energy concentration on feed intake and performance of growing-finishing pig housed individually. *J. Anim. Sci.*, 90, 4362-4372.
- Sauvant D., Perez J.M., Tran G., 2004. Tables of composition and nutritive value of feed materials Pigs, poultry, cattle, sheep, goats, rabbits, horses, fish. INRA Editions, Versailles, France nb pages?
- O'Quinn P.R., Knabe D.A., Gregg E.J., 1997. Efficacy of Natuphos in sorghum-based diets of finishing swine. *J. Anim. Sci.*, 75, 1299-1307.
- van Milgen J., Valancogne A., Dubois S., Dourmad J.Y., Sève B., Noblet J., 2008. InraPorc: A model and decision support tool for the nutrition of growing pigs. *Anim. Feed Sci. Technol.*, 143, 387-405.
- Woyengo T.A., Sands J.S., Guenter W., Nyachoti C.M., 2008. Nutrient digestibility and performance responses of growing pigs fed phytase and xylanase-supplemented wheat-based diets. *J. Anim. Sci.*, 86, 848-857.
- Yáñez J.L., Beltranena E., Cervantes M., Zijlstra R.T., 2011. Effect of phytase and xylanase supplementation or particle size on nutrient digestibility of diets containing distillers dried grains with solubles co fermented from wheat and corn in ileal-cannulated grower pigs. *J. Anim. Sci.*, 89, 113-123.
- Zhang G.J., Yi X.W., Chu L.C., Lu, N., Htoo J., Qiao S.Y., 2011. Effects of Dietary Net Energy Density and Standardized Ileal Digestible Lysine: Net Energy Ratio on the Performance and Carcass Characteristic of Growing-Finishing Pigs Fed Low Crude Protein Supplemented with Crystalline Amino Acids Diets. *Agri. Sci. Ch.* 10, 602-610.