

























- Mallmann F.J.K., dos Santos Rheinheimer D., Ceretta C.A., Cella C., Gomes Minella J.P., Guma R.L., Filipović V., van Oort F., Šimůnek J., 2014. Soil tillage to reduce surface metal contamination – model development and simulations of zinc and copper concentration profiles in a pig slurry-amended soil. *Agric. Ecosyst. Environ.*, 196, 59-68.
- Männer K., 2008. Bioavailability of trace minerals sources in swine. In: P. Schlegel, S. Durosoy, A.W. Jongbloed (Eds), *Trace elements in animal production systems*, 177-186. Wageningen Academic Publishers, Wageningen, The Netherlands.
- Marcato C.E., 2007. Origine, devenir et impact du cuivre et zinc des lisiers porcins. Rôle de la digestion anaérobie. Thèse de doctorat, 184 p.
- Marcato C.E., Pinelli E., Pinelli E., Cecchi M., Winterton P., Guireesse M., 2009. Bioavailability of Cu and Zn in raw and anaerobically digested pig slurry. *Ecotoxicol. Environ. Saf.*, 72, 1538-1544.
- Marszałek M., Kowalski Z., Makara A., 2019. The possibility of contamination of water-soil environment as a result of the use of pig slurry. *Ecol. Chem. Eng. Science*, 26 (2), 313-330.
- Matheri A.N., Belaid M., Seodigeng T., Ngila J.C., 2016. The role of trace elements on anaerobic co-digestion in biogas production. *Proceedings of the world congress on engineering Vol II WCE 2016*, June 29 - July 1, London, U.K.
- McGrath M.C., 1981. Implications of applying copper rich pig slurry to grassland; effects on plant and soil. In : P. L'Hermite, J. Dehandtschutter (Eds), *Copper in animal wastes and sewage sludge*, 144-153. Proc. EEC Workshop, INRA Publisher, Bordeaux, France.
- McGrath S.P., Chaudri A.M., Giller K.E., 1995. Long term effects of metals in sewage sludge on soils, microorganisms and plants. *J. Ind. Microbiol.*, 14, 94-104.
- McGrath S.P., Zhao F.J., Dunham S.J., Crosland A.R., Coleman K., 2000. Long-term changes in the extractability and bioavailability of zinc and cadmium after sludge application. *J. Environ. Qual.*, 29, 875-883.
- Møller H.B., Jensen L.S., Tobiasen L., Hansen M.N., 2007. Heavy metal and phosphorus content of fractions from manure treatment and incineration. *Environ. Technol.*, 28, 1403-1418.
- National Research Council (NRC), 2012. *Nutrient Requirement of Swine*. The National Academies Press, Washington D.C., USA, 389 p.
- Nicholson F.A, Chambers B.J, 2008. Livestock manure management and treatment: implications for heavy metal inputs to agricultural soils. In : P. Schlegel, S. Durosoy, A.W. Jongbloed (Eds), *Trace elements in animal production systems*, 55-62. Wageningen Academic Publishers, Wageningen, The Netherlands.
- Oberleas D., 1973. Phytates. In: *Toxicants occurring naturally in foods*, 363–371. 2nd Ed. National Academies Press, Washington DC.
- O'Dell, 1989. Mineral Interactions Relevant to Nutrient Requirements. *J. Nutr.*, 1832–1838.
- Paboeuf F., Calvar C., Landrain B., Roy H., 2001. Impact de la réduction des niveaux alimentaires en matière azotée totale, en phosphore, en cuivre et en zinc sur les performances et les rejets des porcs charcutiers. *Journées Rech. Porcine*, 33, 49-56.
- Pantelopoulos A., Aronsson H., 2020. Two-stage separation and acidification of pig slurry – Nutrient separation efficiency and agronomical implications. *J. Environ. Manage.*, 280 (111653), 8 p.
- Popovic O., Jensen L.S., 2012. Storage temperature affects distribution of carbon, VFA, ammonia, phosphorus, copper and zinc in raw pig slurry and its separated liquid fraction. *Water Res.*, 46, 3849-3858.
- Popovic O., Hjorth M., Jensen L.S., 2012. Phosphorus, copper and zinc in solid and liquid fractions from full-scale and laboratory-separated pig slurry. *Environ. Technol.*, 33 (18), 2119-2131.
- Poulsen H.D, 1998. Zinc and copper as feed additives, growth factors or unwanted environmental factors. *J. Anim. Feed Sci.*, 7, 135–142.
- Regulation (EU) 2019/4 of the european parliament and of the council of 11 december 2018. <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32019R0004&from=EN>
- Regulation (EU) 2019/6 of the european parliament and of the council of 11 december 2018. <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32019R0006&from=EN>
- Revy P.S., 2003. Diminution des rejets de zinc chez le porc : estimation des besoins et voies d'amélioration de la disponibilité. Thèse de doctorat, 147 p, Agrocampus Ouest.
- Revy P.S., Jondreville C., Dourmad J.Y. and Nys Y., 2004. Assessment of dietary zinc requirement of weaned piglets fed diets with or without microbial phytase. *J. Anim. Physiol. Anim. Nutr.*, 90, 50-59.
- Roméo A., Durosoy S., van Baal J., Bikker P., 2018. Effet de deux sources de cuivre sur les performances et le statut en cuivre de porcelets sevrés. *Journées Rech. Porcine*, 50, 131-136.
- Santos A., Fanguero D., Moral R., Pilar Bernal M., 2018. Composts produced from pig slurry solids: Nutrient efficiency and N-leaching risks in amended soils. *Front. Sustainable Food Syst.*, 2, 12p.
- Schlegel P., 2010. Facteurs de variation de la biodisponibilité du zinc, ajouté sous forme organique ou inorganique, chez deux espèces monogastriques en croissance (poulet et porcelet). Thèse de doctorat, 177 p.
- Schlegel P., Jondreville C., 2011. Interactions phytates - zinc chez le porcelet. *Journées Rech. Porcine*, 43, 95–98.
- Spears J.W., Hansen S.L., 2008. Bioavailability criteria for trace minerals in monogastrics and ruminants. In: P. Schlegel, S. Durosoy, A.W. Jongbloed (Eds), 161–175. *Trace elements in animal production systems*. Wageningen Academic Publishers, Wageningen, The Netherlands.
- Suttle N.F., 2010. *Mineral nutrition of livestock*. 4th Edition. Cabi, Oxfordshire, United Kingdom, 579 p.
- Swinkels J.W.G.M., Kornegay E.T., Verstegen M.W.A., 1994. Biology of zinc and biological value of dietary organic zinc complexes and chelates. *Nutr. Res. Rev.*, 7, 129-149.
- Tessier A., Campbell P.G.C., Bisson M., 1979. Sequential extraction procedure for the speciation of particulate trace metals. *Anal. Chem.*, 51 (7), 844-850.
- Van Heugten E., O'Quinn P.R., Funderburke D.W., Flowers W.L., Spears J.W., 2004. Growth performance, carcass characteristics, plasma minerals, and fecal mineral excretion in grower finisher swine fed diets with levels of trace minerals lower than common industry levels. *J. Swine Health Prod.*, 12 (5), 237-241.
- Villagómez-Estrada S., Pérez J.F., Darwich L., Vidal A., van Kuijk S., Melo-Durán D., Solà-Oriol D., 2020a. Effects of copper and zinc sources and inclusion levels of copper on weanling pig performance and intestinal microbiota. *J. Anim. Sci.*, 98, 1-15.
- Villagómez-Estrada S., Pérez J.F., van Kuijk S., Melo-Durán D., Karimirad R., Solà-Oriol D., 2020b. Effects of two zinc supplementation levels and two zinc and copper sources with different solubility characteristics on the growth performance, carcass characteristics and digestibility of growing-finishing pigs. *J. Anim. Physiol. Anim. Nutr.*, 105, 1–13.
- Yang S., Wen Q., Chen Z., 2020. Impacts of Cu and Zn on the performance, microbial community dynamics and resistance genes variations during mesophilic and thermophilic anaerobic digestion of swine manure. *Bioresour. Technol.*, 312, 11 p.
- Zhang S., Hua Y., Deng L., 2016. Nutrient status and contamination risks from digested pig slurry applied on a vegetable crops field. *Int. J. Environ. Res. Public Health*, 13 (406), 11 p.